



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2016

Ascites secondary to compression of the caudal vena cava by liver abscesses in a cow

Braun, Ueli ; Stettler, M ; Schramm, S ; Hilbe, Monika

Abstract: Background: Ascites due to thrombosis of the caudal vena cava is relatively seldom in cattle. To our knowledge, there have been no reports of ascites secondary to compression of the caudal vena cava by liver abscesses. This case report describes the findings in a 3.7-year-old Brown Swiss cow with this disease. Case presentation: The cow had a distended, pear-shaped abdomen. Ultrasonography of the abdomen showed a massive accumulation of hypoechoic fluid and dilatation of the caudal vena cava. A tentative diagnosis of thrombosis of the caudal vena cava was made, and because of a poor prognosis, the cow was slaughtered. Incision of the abdominal wall released approximately 100 litres of yellow, clear, non-malodorous fluid from the peritoneal cavity. The liver was markedly enlarged, and two abscesses, approximately 10 cm in diameter, were situated immediately adjacent to the caudal vena cava causing compression of the vein. No microbiology culture was done. The final diagnosis was ascites secondary to compression of the caudal vena cava by two liver abscesses and secondary hepatic fibrosis. Conclusion: This report suggests that compression of the caudal vena cava by liver abscesses should be included in the list of differential diagnosis of conditions associated with ascites and dilatation of the caudal vena in cattle.

DOI: <https://doi.org/10.15744/2348-9790.4.105>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-125641>

Journal Article

Accepted Version

Originally published at:

Braun, Ueli; Stettler, M; Schramm, S; Hilbe, Monika (2016). Ascites secondary to compression of the caudal vena cava by liver abscesses in a cow. *Journal of Veterinary Science and Animal Husbandry*, 4(1):1-7.

DOI: <https://doi.org/10.15744/2348-9790.4.105>

1 Ascites secondary to compression of the caudal vena cava by liver
2 abscesses in a cow

3
4 Ueli Braun^{§1}, Manuela Stettler¹, Stephanie Schramm¹, Monika Hilbe²

5
6 ¹Department of Farm Animals, Vetsuisse Faculty, University of Zurich

7 ²Institute of Veterinary Pathology, Vetsuisse Faculty, University of Zurich

8
9 [§]Corresponding author

Abstract

Background

Ascites due to thrombosis of the caudal vena cava is relatively seldom in cattle. To our knowledge, there have been no reports of ascites secondary to compression of the caudal vena cava by liver abscesses. This case report describes the findings in a 3.7-year-old Brown Swiss cow with this disease.

Case presentation

The cow had a distended, pear-shaped abdomen. Ultrasonography of the abdomen showed a massive accumulation of hypoechoic fluid and dilatation of the caudal vena cava. A tentative diagnosis of thrombosis of the caudal vena cava was made, and because of a poor prognosis, the cow was slaughtered. Incision of the abdominal wall released approximately 100 litres of yellow, clear, non-malodorous fluid from the peritoneal cavity. The liver was markedly enlarged, and two abscesses, approximately 10 cm in diameter, were situated immediately adjacent to the caudal vena cava causing compression of the vein. **No microbiology culture was done.** The final diagnosis was ascites secondary to compression of the caudal vena cava by two liver abscesses **and secondary hepatic fibrosis.**

Conclusions

This report suggests that compression of the caudal vena cava by liver abscesses should be included in the list of differential diagnosis of conditions associated with ascites and dilatation of the caudal vena in cattle.

Keywords: Cattle, ascites, liver abscess, caudal vena cava, compression

Background

Ascites is excessive accumulation of fluid in the peritoneal cavity. The ascites fluid can be characterised as inflammatory, non-inflammatory, chylous, urine-like, haemorrhagic or bilious [1]. In

cattle, ascites fluid is usually non-inflammatory and associated with vascular congestion due to chronic right-sided heart failure, a mediastinal mass, chronic liver or kidney disease, ileus of the small intestines, enteropathies, peritoneal tumours or thrombosis of the caudal vena cava [1]. There are several reports on ascites associated with thrombosis of the caudal vena cava in cattle [2-5], but to the authors' knowledge, there are no reports on ascites secondary to compression of this vein. This case report describes the clinical, ultrasonographic and pathological findings in a 3.7-year-old Brown Swiss cow, which was five months pregnant and had ascites secondary to compression of the caudal vena cava by liver abscesses.

Case presentation

For several days before referral to our clinic, the cow had a marked decrease in appetite and faecal output and progressive distension of the abdomen. On admission, the cow was in poor general health, had a marked decrease in appetite and a pear-shaped abdomen (Fig. 1). The rectal temperature was 38.5 °C, the heart rate 68 bpm and the respiratory rate 28 breaths per min. Rumen motility was severely decreased, the rumen was relatively full and there was no distinct layering of rumen contents. Tests for reticular foreign bodies and swinging ~~and auscultation~~ and percussion ~~and~~ auscultation on both sides of the abdomen were negative. Intestinal motility was markedly decreased, and the cow passed small amounts of dark, watery, malodorous faeces. The results of a urine test strip (Combur⁹-Test, Roche, Basel) were unremarkable. An increased volume of peritoneal fluid was suspected based on sloshing felt during transrectal palpation.

The results of haematological and serum biochemical analyses showed haemoconcentration (~~haematocrit 38 %, reference interval 33—36 %~~), increase in γ -GT activity (~~56 U/l, reference interval 15—25 U/l~~), azotaemia (~~urea 7.7 mmol/l, reference interval 3.5—6.5 mmol/l~~), hypocalcaemia (~~calcium 2.1 mmol/l, reference interval 2.3—2.6 mmol/l~~) and metabolic acidosis (Table 1) (~~venous~~

58 ~~blood pH 7.36, bicarbonate 21.1 mmol/l, base excess -3.1 mmol/l; reference interval blood pH 7.40–~~
59 ~~7.50, bicarbonate 20–30 mmol/l, base excess -2 to +2 mmol/l).~~

60 Transcutaneous ultrasonography was performed using a 5.0 MHz convex transducer. It showed a
61 massive amount of hypoechoic fluid in the peritoneal cavity, and the reticulum (Fig. 2), intestines
62 (Fig. 3) and liver (Fig. 4) were displaced dorsally by the ascites fluid. The liver parenchyma appeared
63 heterogeneous and hyperdense on ultrasonograms ~~because of congestion~~. The caudal vena cava was
64 circular and dilated in cross-section (Fig. 5) and did not have a normal triangular shape [6]. The
65 gallbladder wall was thicker than normal because of oedema (Fig. 4). Ultrasound-guided
66 abdominocentesis yielded yellow, clear, non-malodorous fluid with a specific gravity of 1.034
67 (reference interval < 1.015), a protein concentration of 46 g/l (reference interval < 30 g/l), urea
68 concentration of 7.5 mmol/l (reference interval similar to that of blood) and creatinine concentration
69 of 94.3 µmol/l (reference interval similar to that of blood). Cytological evaluation of the abdominal
70 fluid showed a few erythrocytes, leukocytes and macrophages, and was classified as a modified
71 transudate. Based on all the findings, a tentative diagnosis of non-inflammatory ascites secondary to
72 thrombosis of the caudal vena cava was made. The cow was slaughtered because of a poor prognosis,
73 and a postmortem examination was carried out. Approximately 100 litres of clear, non-malodorous
74 fluid escaped from the abdomen when the abdominal wall was incised (Fig. 6). The liver was
75 markedly increased in size and firmer than normal and had a bumpy surface. Two abscesses with a
76 diameter of approximately 10 cm were located immediately adjacent to the caudal vena cava and
77 compressed it (Fig. 7, 8). There was no thrombus in the caudal vena cava (Fig. 9). Histopathology of
78 the liver parenchyma revealed a diffuse fibrosis, which was especially severe around the centrilobular
79 veins and was often forming fibrous bridges between the periportal spaces (bridging fibrosis).
80 Hepatocytes and hepatic lobules were severely atrophic and midzonal sinusoids were heavily
81 congested and dilated (severe hepatic chronic congestion). Portal tracts were characterized by severe
82 bile duct proliferation and multiplication of arterioles. Several vessel walls were moderately thickened

83 and centrolobular vein. The final diagnosis was ascites attributable to compression of the caudal vena
84 cava by two liver abscesses and secondary hepatic fibrosis.

85 The causes of ascites and the diagnostic work-up of cattle with ascites were recently described in
86 detail [1]. Ultrasonography, abdominocentesis and examination of peritoneal fluid are critical
87 components for determining the aetiology of the problem. Inflammatory and non-inflammatory
88 conditions can be differentiated by peritoneal fluid analysis. Uroperitoneum is diagnosed when the
89 creatinine concentration of peritoneal fluid is higher than that of serum, and an increased
90 concentration of bilirubin in peritoneal fluid usually indicates bile peritonitis. Rupture of a tumour or
91 blood vessel results in haemorrhagic peritoneal fluid, and peritoneal fluid that appears milky and has a
92 high lymphocyte count may indicate chyloabdomen. The most important finding in this case study,
93 other than the massive accumulation of peritoneal fluid, was the change in shape of the caudal vena
94 cava in cross-section. The caudal vena cava normally appears triangular in cross-section [6], but
95 becomes oval to circular with congestion of the vein. Changes in the shape of the vein are of greater
96 diagnostic importance than an increase in the diameter of the vein [7]. Congestion of the caudal vena
97 cava occurs with right-sided heart failure and with obstruction or compression of the vein. In cattle
98 with right-sided heart failure, there is also congestion of the jugular veins, which did not occur in the
99 present case. Our tentative diagnosis of thrombosis of the caudal vena cava appeared to be the most
100 plausible cause because it occurs relatively often in cattle [2-4,7-9]. However, in these cases, a
101 thrombus can rarely be seen via ultrasonography [8-10] because most are located in the part of the
102 caudal vena cava that is obscured by the lungs. An outstanding yet time-consuming method for
103 identifying a thrombus in the caudal vena cava is intraabdominal ultrasonography of the vein through
104 a right-flank laparotomy [11]. Our tentative diagnosis was proved incorrect during postmortem
105 examination, which revealed compression of the caudal vena cava by two liver abscesses.

106 Unfortunately, ultrasonography could not be used to visualize the abscesses because they were located
107 in a part of the liver obscured by the lungs. It is highly likely that a correct diagnosis would have been

possible using intraabdominal ultrasonography [11]. The liver abscesses were immediately adjacent to the caudal vena cava and thus led to compression and subsequent congestion of the vein. Liver abscesses are the most common cause of thrombosis of the caudal vena cava in cattle [12-18] and, similar to our case, are situated immediately adjacent to the vein. However, liver abscesses usually rupture into the caudal vena cava and form a fibrinous thrombus attached to the intima, which partially or completely obstructs the lumen of the vein. This did not occur in our case.

Severe ascites usually is a sequel to increased hydrostatic and decreased oncotic pressure. In the case described in this report, increased hydrostatic pressure in the caudal vena cava was caused by an abscess that compressed the vein and impaired blood flow to the heart, and the decreased oncotic pressure was the result of liver fibrosis caused by congestion.

Conclusion

In conclusion, the change in the cross-sectional shape of the caudal vena cava from triangular to circular or oval on ultrasonography and the features found through the necropsy suggest that the ascites was caused by compression performed by the abscess and the secondary hepatic fibrosis.

~~A change in the cross-sectional shape of the caudal vena cava from triangular to circular or oval on ultrasonograms is abnormal in cattle. In such cases, right-sided heart failure, thrombosis of the caudal vena cava and compression of the caudal vena cava by an abscess must be part of the differential diagnosis.~~

Competing interests

The authors declare that they have no competing interests.

Author's contributions

UB prepared the manuscript and supervised the clinical and ultrasonographic examination, MS and SS examined the heifer, postmortem examination was performed by MH. All authors have read and approved the manuscript.

Authors' information

¹Department of Farm Animals, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland

²Institute of Veterinary Pathology, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland.

Acknowledgements

The authors thank Dr Rainer Nussbaumer for referring the cow to our clinic, the technicians of the Medical Laboratory for the haematological and biochemical analyses and the animal assistants for looking after the cow.

References

1. Braun U. Ascites in cattle – Ultrasonographic findings and diagnosis. *Vet Clin North Am Food Anim Pract.* 2016;32:In press (March 2016).
2. Braun U, Schefer U, Gerber D, Föhn J. Ultrasonographic findings in a cow with ascites due to thrombosis of the caudal vena cava. *Schweiz Arch Tierheilk.* 1992;134:235-41.
3. Braun U, Schweizer G, Wehbrink D, Müller R, Hilbe M. Ultraschallbefunde bei einem Rind mit Aszites infolge Thrombose der Vena cava caudalis. *Tierärztl Prax.* 2005;33(G):389-94.
4. Braun U, Feller B, Trachsel D, Rütten M, Augsburg H, Riond B, Sydler T. Lungenabszess, Pleuraerguss und Aszites bei einem Rind mit Vena-cava-caudalis-Thrombose. *Dtsch tierärztl Wschr.* 2007;114:165-70.
5. Braun U. Clinical findings and diagnosis of thrombosis of the caudal vena cava in cattle. *Vet J.* 2008;175:118-25.
6. Braun U. Ultrasonography of the liver in cattle. *Vet Clin North Am Food Anim Pract.* 2009;25:591-609.
7. Braun U, Flückiger M, Feige K, Pospischil A. Diagnosis by ultrasonography of congestion of the caudal vena cava secondary to thrombosis in 12 cows. *Vet Rec.* 2002;150:209-13.
8. Braun U, Salis F, Gerspach C. Sonographischer Nachweis eines echogenen Thrombus in der Vena cava caudalis bei einer Kuh. *Schweiz Arch Tierheilk.* 2003;145:340-1.

- 157 9. Mohamed T, Sato H, Kurosawa T, Oikawa S. Ultrasonographic localisation of thrombi in the
158 caudal vena cava and hepatic veins in a heifer. *Vet J.* 2004;168:103-6.
- 159 10. Simpson KM, Streeter RN, Cramer S, Lamm CG, Love BC. Caudal vena caval thrombosis
160 following treatment of deep digital sepsis. *Can Vet J.* 2012;53:182-6.
- 161 11. Sigrist I, Francoz D, Leclère M, Buczinski S. Antemortem diagnosis of caudal vena cava
162 thrombosis in 2 cows. *J Vet Intern Med.* 2008;22:684-6.
- 163 12. Rubarth S. Hepatic and subphrenic abscesses in cattle with rupture into vena cava caudalis. *Acta*
164 *Vet Scand.* 1960;1:363-82.
- 165 13. Stöber M. Pyogene Thrombosen der Vena cava caudalis beim Rind. *Schweiz Arch Tierheilk.*
166 1966;108:613-21.
- 167 14. Selman IE, Wiseman A, Petrie L, Pirie HM, Breeze RG. A respiratory syndrome in cattle
168 resulting from thrombosis of the posterior vena cava. *Vet Rec.* 1974;94:459-66.
- 169 15. Rebhun WC, Rendano VT, Dill SG, King JM, Pearson EG. Caudal vena caval thrombosis in four
170 cattle with acute dyspnea. *J Am Vet Med Assoc.* 1980;176:1366-9.
- 171 16. Ikawa H, Narushima T, Kohno T. Bacteriology of caudal vena caval thrombosis in slaughter
172 cattle. *Vet Rec.* 1987;120:184-6.
- 173 17. Mills LL, Pace LW. Caudal vena caval thrombosis in a cow. *J Am Vet Med Assoc.*
174 1990;196:1294-6.
- 175 18. Radostits OM, Gay CC, Hinchliff KW, Constable PD. Caudal vena caval thrombosis (posterior
176 vena caval thrombosis) and embolic pneumonia in cattle. In: *Veterinary Medicine. A Textbook of*
177 *the Diseases of Cattle, Horses, Sheep, Pigs, and Goats.* In: Radostits OM, Gay CC, Hinchliff
178 KW, Constable PD, editors. Edinburgh: Saunders Elsevier; 2007. p. 516-7.

179 Figure legends

180 **Figure 1 - Pear-shaped abdomen in a cow with ascites.** Photo showing a pear-shaped abdomen
181 in a Brown-Swiss cow with ascites due to compression of the caudal vena cava by two liver
182 abscesses.

183 **Figure 2 - Ultrasonogram of the reticular region.** Ultrasonogram of the reticular region of a cow
184 with ascites due to compression of the caudal vena cava by two liver abscesses, imaged from the
185 sternal part of the ventral abdomen. The reticulum is displaced dorsally by anechoic ascites fluid. 1
186 Ventral abdominal wall, 2 Musculophrenic vein, 3 Diaphragm, 4 Ascites fluid, 5 Reticulum, 6 Anterior
187 dorsal blind sac of the rumen, Cr Cranial, Cd Caudal.

188 **Figure 3 - Ultrasonogram of ascites.** Ultrasonogram of ascites in a cow with compression of the
189 caudal vena cava by two liver abscesses, imaged from the right flank. Loops of small intestines are
190 displaced dorsally by anechoic ascites fluid. 1 Lateral abdominal wall, 2 Extraomental ascites fluid, 3
191 Greater omentum, 4 Intraomental ascites fluid, 5 Small intestines, Ds Dorsal, Vt Ventral.

192 **Figure 4 - Ultrasonogram of the liver and gallbladder.** Ultrasonogram of the liver and gallbladder
193 of a cow with ascites due to compression of the caudal vena cava by two liver abscesses, imaged
194 from the 10th intercostal space on the right side. The liver and the gallbladder are displaced from
195 the abdominal wall by anechoic ascites fluid. The wall of the gallbladder appears thicker than normal
196 is thickened because of oedema (congestion). 1 Lateral abdominal wall, 2 Ascites fluid, 3 Liver, 4
197 Gallbladder, Ds Dorsal, Vt Ventral.

198 **Figure 5 - Ultrasonogram of the caudal vena cava.** Ultrasonogram of the caudal vena cava in a
199 cow with compression of the caudal vena cava by two liver abscesses, imaged from the 11th
200 intercostal space on the right side. The caudal vena cava, which normally is triangular in cross-
201 section, is round because of congestion and dilation. 1 Lateral abdominal wall, 2 Liver, 3 Hepatic
202 vein, 4 Caudal vena cava, 5 Right hepatic vein, Ds Dorsal, Vt Ventral.

203 **Figure 6 - Yellow ascites fluid escapes from the peritoneal cavity.** A large volume of yellow
204 ascites fluid escapes from the peritoneal cavity after incision of the abdominal wall during

205 postmortem examination of a Brown-Swiss cow with compression of the caudal vena cava by two
206 liver abscesses.

207 **Figure 7 - Liver with two abscesses.** The diaphragmatic fascia of the liver of a Brown-Swiss cow
208 with two abscesses (arrows) in the upper margin of the liver and immediately adjacent to the caudal
209 vena cava. The liver is whitish-grey and enlarged because of congestion of the caudal vena cava.

210 **Figure 8 - Pus oozes from an incised liver abscess.** Yellow viscous pus (arrow) oozes from an
211 incised liver abscess that was compressing the caudal vena cava in a Brown-Swiss cow.

212 **Figure 9 - Opened caudal vena.** The caudal vena cava (arrow) of a Brown-Swiss cow with
213 compression of the vein by two liver abscesses. The incised and opened caudal vena cava has no
214 macroscopic abnormalities, such as a thrombus.

215 Table 1: Selected haematological and biochemical results and venous blood gas analysis in a cow
216 with ascites secondary to compression of the caudal vena cava by liver abscesses

Variable	Cow	Normal range
Haematocrit (%)	38	33 – 36
γ -GT (U/l)	56	15 – 25
Urea (mmol/l)	7.7	3.5 – 6.5
Calcium (mmol/l)	2.1	2.3 – 2.6
Venous bood pH	7.36	7.40 – 7.50
Bicarbonate (mmol/l)	21.1	20 – 30
Base excess (mmol/l)	-3.1	-2 to +2